



A service of the National Library of Medicine  
and the National Institutes of Health

www.pubmed.gov

My NCBI

[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

1: Invest Ophthalmol Vis Sci. 1998 Jul;39(8):1346-51.

Links

### Reoxygenation injury in a cultured corneal epithelial cell line protected by the uptake of lactoferrin.

Shimmura S, Shimoyama M, Hojo M, Urayama K, Tsubota K.

Department of Ophthalmology, Tokyo Dental College, Chiba, Japan.

**PURPOSE:** To investigate whether reoxygenation after extended hypoxia causes cellular damage in cultured corneal epithelial cells and to demonstrate the protective effects of lactoferrin. **METHODS:** Immortalized human corneal epithelial cells (T-HCECs) were cultured to confluence in 96-well culture plates, subjected to stringent hypoxia (1% O<sub>2</sub>, 5% CO<sub>2</sub>, 94% N<sub>2</sub> at 37 degrees C) for 24 hours, and returned to normoxic conditions (5% CO<sub>2</sub>, 95% air at 37 degrees C). Cell viability was observed by 1 microM propidium iodide staining 0, 2, 4, and 6 hours after reoxygenation. Inhibition studies were performed after 2 hours' reoxygenation, using 2 mM iron chelator desferrioxamine and 0.2 mg/ml lactoferrin. Confocal immunocytochemistry for human lactoferrin and western blot analysis for lactoferrin-induced ferritin were performed in cultured T-HCECs to demonstrate the internalization of lactoferrin after application. **RESULTS:** After 2 hours, reoxygenation of T-HCECs after hypoxia produced an increase in cell death that was significantly greater than that observed in normoxic control cells or in cells subjected to hypoxia for the same time span without reoxygenation. The addition of desferrioxamine and lactoferrin at the time of reoxygenation significantly attenuated cellular damage. Confocal immunocytochemistry revealed that lactoferrin is taken into the cytoplasm of T-HCECs as early as 30 minutes after application. This was also demonstrated in western blot analysis by the upregulation of intracellular ferritin at 18 hours by the addition of iron-bound lactoferrin but not by iron-free lactoferrin. **CONCLUSION:** Reoxygenation is responsible for increased cellular damage after extensive hypoxia, which is attenuated by chelators of free iron in the cytosol, including the major tear protein lactoferrin.

PMID: 9660482 [PubMed - indexed for MEDLINE]

### Related Links

Human corneal epithelial cell viability and morphology after dilute alcohol exposure. [Ophthalmol Vis Sci. 2007]

Defensin expression by the cornea: multiple signalling pathways mediate IL-1beta stimulation of hBD-2 expression by human corneal epithelial cells. [Ophthalmol Vis Sci. 2003]

The effect of hypoxia/reoxygenation on the cellular function of intestinal epithelial cells. [J Trauma. 1999]

Reoxygenation results in cell death of human alveolar epithelial cells. [Am J Physiol Lung Cell Mol Physiol. 1994]

Reactive oxygen species and rat renal epithelial cells during hypoxia and reoxygenation. [Kidney Int. 1991]

See all Related Articles...

Display AbstractPlus Show 20 Sort by Send to